

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 627 217 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
06.08.2003 Bulletin 2003/32

(51) Int Cl.7: **A61K 7/50, A61K 7/06**

(21) Application number: **94107637.4**

(22) Date of filing: **17.05.1994**

(54) **Stable conditioning shampoo having a high foam level containing a silicone conditioner, a cationic quaternary acrylate copolymer, an anionic surfactant and polyethyleneimine**

Stabile konditionierendes Haarpflegershampoo mit hoher Schaumkraft enthaltend ein Silicone Konditionierungsmittel, ein Kationisches quaternaires Acrylat-Kopolymer, ein anionisches Tensid und Polyethyleneimine

Shampooing de conditionnement stable ayant une action moussante élevée contenant un conditionneur silicone, un copolymère acrylate cationique quaternaire, un tensioactive anionique et polyethyleneimine

(84) Designated Contracting States:
DE DK ES FR GB IT SE

(30) Priority: **17.05.1993 US 62606**

(43) Date of publication of application:
07.12.1994 Bulletin 1994/49

(73) Proprietors:
• **UNILEVER PLC**
London EC4P 4BQ (GB)
Designated Contracting States:
GB
• **UNILEVER N.V.**
3013 AL Rotterdam (NL)
Designated Contracting States:
DE DK ES FR IT SE

(72) Inventors:

- **Janchitraponvej, Ben**
Niles, IL (US)
- **Brown, William J.**
Flossmoor, IL (US)

(74) Representative: **Griffiths, Helen Sarah et al**
Unilever PLC,
Patent Division,
Colworth House
Sharnbrook, Bedford MK44 1LQ (GB)

(56) References cited:
WO-A-94/06409 **GB-A- 1 195 158**

EP 0 627 217 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

THIS PAGE BLANK (USPTO)

Description

FIELD OF THE INVENTION

5 [0001] The present invention is directed to a hair conditioning shampoo composition and to a method of treating hair with the composition to provide the hair with improved wet stage and dry stage conditioning properties as well as other conditioning properties, such as softness, without residual build-up of conditioning agents on the hair. The conditioning shampoo also thoroughly cleanses the hair, while conditioning, with a cleansing detergent that develops an unexpectedly high foam level and unexpected stability containing an anionic detergent, a protonated (cationic) polyethyleneimine

10 conditioning agent and, optionally, one or more silicone conditioning agents. More particularly, the present invention is directed to a hair conditioning shampoo composition including one or more anionic cleansing surfactants, such as ammonium lauryl sulfate or ammonium lauryl ether sulfate (ALES), a protonated polyethyleneimine, and an acrylate/acrylamide copolymer conditioning agent that is stable over extended periods of time at elevated temperature.

15 [0002] Surprisingly, the composition develops copious amounts of foam for a conditioning shampoo while achieving excellent conditioning benefits, and is unexpectedly stable while having a combination of a strong anionic cleansing detergent, such as a long chain alkyl sulfate, or a partially ethoxylated long chain alkyl sulfate or sulfonate together with a cationic polyethyleneimine and a silicone conditioning agent. The anionic surfactant/cationic polyethyleneimine conditioning agent components are compatible and stable while developing surprisingly high amounts of foam without the problem of anionic surfactant-cationic conditioning agent incompatibility.

BACKGROUND OF THE INVENTION

20 [0003] Soiled human hair is shampooed to remove sebum that is naturally secreted by the head as well as soil and other atmospheric contaminants that accumulate on the hair. Sebum, in particular, accumulates on the hair in a relatively short period of time leaving the hair with a greasy, dirty feel and poor manageability. The most effective shampoos for cleansing the hair for removal of the atmospheric contaminants and sebum, are those that contain high lather synthetic anionic detergents, such as the long chain alkyl sulfates, the partially ethoxylated long chain alkyl sulfates and the long chain sulfonates. These synthetic anionic detergents are very effective for cleansing the hair but, after rinsing with water, leave the hair with a dried touch, usually called "creak" and result in hair, when wet, that is in an extremely tangled and unmanageable after-shampoo condition.

30 [0004] Thoroughly cleansed hair is extremely difficult to comb, in either the wet or dry state because the individual hair fibers tend to snarl, kink, and interlock with each other. Particularly, prior to complete drying of thoroughly cleansed hair, in this after-shampoo stage, the hair is very difficult to comb or brush. Even after complete drying, the thoroughly cleansed hair remains difficult to comb or brush and does not set well. Thoroughly clean, dried hair also has undesirable electrostatic properties in a low humidity atmosphere that causes the hair to "fly away", thereby further reducing the combing or brushing property of the hair. Generally, these above-outlined problems that result from synthetic detergent cleansing of the hair, particularly in the high-lather synthetic anionic detergents, have been alleviated either by the after-shampoo treatment of the hair with hair conditioners, for example, in the form of a hair rinse, or by including hair conditioners directly within the shampoo composition.

40 [0005] After-shampoo hair conditioning compositions are easily formulated but are inconvenient to use because of the necessity of applying the conditioner to the hair in a separate stage, after shampooing. The preparation of a conditioning shampoo has been more difficult because of inherent incompatibility problems between anionic surfactants and the cationic compounds that are good conditioning agents. Contact between an anionic surfactant and a cationic surfactant or cationic polymer produces a precipitate that forms immediately or causes an interaction between the anionic and cationic compounds that significantly reduces their respective cleaning and conditioning properties, and especially a very noticeable severe loss of foam attributed by the anionic cleansing surfactant. The reduction in cleansing and conditioning effectiveness is observed even in compositions wherein the anionic and cationic compounds do not precipitate from the composition but remain in solution or suspension. This incompatibility between an anionic surfactant and a cationic conditioning compound is well recognized by those skilled in the art. For example, Sagarin in *Cosmetics*, Interscience Publishers, Inc., New York, p. 538 (1957), states that anionic and cationic compounds cannot be used in combination because they react to form insoluble salts.

50 [0006] A partial solution to this incompatibility problem in the formulation of conditioning shampoos is exemplified by the following patents that disclose compositions that contain surfactants that are not anionic, e.g., non-ionics, amphoterics and zwitterionics together with cationic conditioning compounds: U.S. Patent No. 3,849,348 to Hewitt; U.S. Patent No. 3,990,991 to Gerstein; and U.S. Patent No. 3,822,312 to Sato.

55 [0007] Another problem inherent in formulating a conditioning shampoo is an instability problem that results when water-insoluble conditioning agents are also included in the conditioning shampoo composition, such as the non-volatile silicones that are well recognized in the art as providing a degree of softness to the hair.

[0008] Silicones in shampoo compositions have been disclosed in a number of different patents: U.S. Patent No. 2,826,551, March 11, 1958 to Green; U.S. Patent No. 3,964,500, June 22, 1976 to Drakoff; U.S. Patent No. 4,364,837, December 21, 1982 to Pader; British Patent No. 849,433, September 28, 1960 to Woolston; U.S. Patent No. 4,741,855 to Grote, et al.; U.S. Patent Nos. 4,788,006 and 4,902,499 to Bolich, Jr., et al. and U.S. Patent No. 4,704,272 to Oh, et al. The silicones are well known to substantially reduce the foaming of anionic cleansing surfactants.

[0009] A particularly difficult problem to solve in silicone-containing conditioning shampoos is that of providing a conditioning shampoo that provides excellent cleansing of the hair while providing high foaming and, at the same time, also has excellent conditioning performance. The capability of providing excellent conditioning, cleansing and foam levels is achieved with the compositions of the present invention. In accordance with one embodiment of the present invention, it has been found that a combination of (1) a cationic (protonated) polyethyleneimine (PEI) in an amount of about 0.01% to about 4% by weight of the composition, preferably about 0.01% to about 1% by weight, together with (2) a cationic oil-soluble, water-dispersible cross-linked quaternary acrylate/acrylamide copolymer (Polyquaternium 32), in an amount of about 0.1% to about 20% based on the total weight of the composition, is stable over long periods of time while providing excellent simultaneous conditioning, cleansing and foaming. Optionally, a number of materials can be included in the conditioning shampoos of the present invention for purposes of thickening and stabilization such as xanthan gum, long chain acyl derivatives, long chain amide oxides or amine oxides, and long chain alkanolamides, as disclosed in U.S. Patent Nos. 4,788,006; 4,704,272; and 4,741,855, hereby incorporated by reference.

SUMMARY OF THE INVENTION

[0010] In accordance with the principles of the present invention, it has been found, surprisingly, that a conditioning shampoo containing an anionic cleansing surfactant, in an amount of about 5% to about 65% by weight, preferably about 5% to about 25% by weight, e.g., (C₁₂-C₂₂) alkyl sulfate, and/or a long chain (C₁₂-C₂₂) partially or fully ethoxylated alkyl sulfate, and/or a long chain (C₁₂-C₂₂) alkyl sulfonate, or alkyl ether sulfonate, and a combination of cationic polymers comprising (1) a cationic (protonated) polyethyleneimine in an amount of about 0.1% to about 4% by weight, preferably about 0.1% to about 1% by weight, and (2) a cationic oil-soluble, water-dispersible, cross-linked quaternary acrylate/acrylamide copolymer (Polyquaternium 32), in an amount of about 0.1% to about 20% by weight of the composition, provides excellent foaming, conditioning and stability, without settling of water-insoluble materials due to anionic-cationic complexing. The composition preferably also includes one or more silicone conditioning agents, and has extended product stability, excellent overall conditioning to human hair, particularly superior wet and dry combing properties, and unexpectedly maintains very high levels of foam.

[0011] The compositions of the present invention are stable and do not exhibit the inherent anionic surfactant/cationic polymer, (polyethyleneimine and/or Polyquaternium 32 conditioning agents) incompatibility while providing excellent cleansing, conditioning and foam levels in a conditioning shampoo. It was further surprisingly and unexpectedly found that hair treated with the compositions of the present invention is thoroughly cleansed at high foam levels and exhibits improved physical and cosmetic properties, such as gloss, wet combing, dry combing, thickness, manageability, softness and body.

[0012] Therefore, an aspect of the present invention is to provide a hair-treating composition that cleanses the hair and imparts improved physical properties and cosmetic properties to the hair in a single application from a mild conditioning shampoo that develops unexpectedly high foam quantities.

[0013] Another aspect of the present invention is to provide a physically stable conditioning shampoo containing an anionic surfactant, and a combination of cationic polymers that provide hair conditioning and composition stability, wherein the composition develops excellent foam levels and can be formulated at room temperature.

[0014] Another aspect of the present invention is to provide a new and improved conditioning shampoo containing a strong anionic detergent, such as a long chain alkyl sulfate, long chain alkyl ether sulfate, and/or long chain sulfonate, that is compatible with cationic conditioning agents, and that maintains an unexpectedly high foam level although the composition contains a silicone conditioning agent.

[0015] Still another aspect of the present invention is to provide a new and improved conditioning shampoo including about 5% to about 25% of an anionic surfactant; about 0.01% to about 4% of a cationic polyethyleneimine that surprisingly provides composition stability and added conditioning benefits, particularly increased wet and dry combing and reduced static (fly away) for better manageability; and about 0.1% to about 20% of a cationic polymeric conditioning agent (Polyquaternium 32), about 0.01% to about 10% of a silicone conditioning agent, and optionally any known emulsion stabilizer and/or a viscosity increasing agent for added stability of aqueous emulsions, each in an amount of about 0.1% to about 10% by weight, active, preferably about 0.1% to about 5% by weight, capable of raising the viscosity to at least 3 Pa.s.

[0016] A further aspect of the present invention is to provide a new and improved method of cleansing and conditioning hair, simultaneously, with a composition containing one or more anionic surfactants; a cationic (protonated) polyethyleneimine having an increased charge density; a cationic acrylate/acrylamide copolymer conditioning agent;

and one or more silicone conditioning agents while providing high foam levels, excellent cleansing, excellent conditioning in a stable conditioning shampoo and is capable of being mixed together at room temperature.

[0017] Still another aspect of the present invention is to provide a new and improved conditioning shampoo having a pH in the range of about 4 to about 7, preferably about 5 to about 6, including about 5% to about 65% of an anionic surfactant; polyethyleneimine in an amount of about 0.01% to about 4%, preferably about 0.01% to about 1% by weight; optionally about 0.1% to about 20% of a cationic, nitrogen-containing acrylate/acrylamide copolymer conditioning agent; and optionally about 0.5% to about 10% of a cationic silicone conditioning agent.

[0018] Another object of the present invention is to provide a new and improved conditioning shampoo having a pH in the range of about 4 to about 7, preferably about 5 to about 6, including about 5% to about 65% of an anionic surfactant; protonated polyethyleneimine in an amount of about 0.01% to about 4%, preferably about 0.01% to about 1% by weight, and having a cationic polymer charge density of at least about 10 milliequivalents per gram, preferably about 15 to about 20 milliequivalents per gram; about 0.1% to about 20% of a cationic acrylate/acrylamide copolymer (Polyquaternium 32) conditioning agent; and about 0.5% to about 10% of a non-volatile silicone material.

[0019] The above and other aspects and advantages of the present invention will become apparent from the following detailed description of the preferred embodiments.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0020] The aqueous conditioning shampoo compositions of the present invention generally include water in an amount of about 60% to about 80-90% by weight; an anionic surfactant preferably in an amount of about 5% to about 25% by weight of the composition; a polyethyleneimine in an amount of about 0.01% to about 4%, preferably about 0.01% to about 1% by weight; optionally a silicone conditioning agent in an amount of about 0.1% to about 10% by weight of the composition; and Polyquaternium 32, a cationic conditioning agent, in an amount of about 0.1% to about 20% by weight of the composition.

[0021] The conditioning shampoo of the present invention provides the hair with improved physical and cosmetic conditioning properties, such as gloss, thickness, softness, and manageability, including excellent wet and dry combing properties and body simultaneously with excellent cleansing at high foam levels in a mild conditioning shampoo. As will be demonstrated more fully hereinafter, it is surprising and unexpected that the composition of the present invention, including an anionic cleansing detergent, and a cationic conditioning compound is able to provide the demonstrated cleansing at such a high foam level in a stable composition containing a silicone conditioning agent.

[0022] The anionic cleansing surfactant used in the composition and method of the present invention can be any of the anionic surfactants known or previously used in the art of hair shampoos. An anionic cleansing surfactant should be included in the composition of the present invention to effectively cleanse the hair and generates a high, stable foam level that consumers equate with cleaning efficiency. While nonionic and amphoteric surfactants have not been as effective in cleansing the hair and do not provide the high foam level desired by consumers, surprisingly, it has been found that the composition of the present invention provides excellent foam levels with the less strong anionic cleansing detergents or with the strong anionic detergents at levels generally below about 9% by weight of the composition, particularly when the foam level is boosted with one or more common foam boosters, such as a betaine or other foam booster. Optionally, nonionic amphoteric and/or zwitterionic surfactants can be included in the compositions of the present invention in addition to one or more anionic surfactants, to help stabilize foam, to provide a suitable viscosity, or to give other functional or esthetic properties to the composition.

[0023] Usually, the anionic cleansing surfactant includes a hydrophobic moiety, such as a carbon chain including from about eight carbon atoms to about 30 carbon atoms, and particularly from about 12 carbon atoms to about 22 carbon atoms and further includes a hydrophilic moiety, such as a sulfate, sulfonate, carbonate, phosphate or carboxylate. Often, the hydrophobic carbon chain is etherified, such as with ethylene oxide or propylene oxide, to impart a particular physical property, such as increased water-solubility or reduced surface tension, to the anionic cleansing surfactant.

[0024] Suitable anionic cleansing surfactants include, but are not limited to, compounds in the classes known as alkyl sulfates, alkyl ether sulfates, alkyl sulfonates, alkyl ether sulfonates, sulfate esters of an alkylphenoxy polyoxyethylene ethanol, alpha-olefin sulfonates, beta alkylalkoxy alkene sulfonates, alkyl arylsulfonates, alkyl carbonates, alkyl ether carboxylates, fatty acids, succinamates, sulfosuccinates, sarcosinates, octoxynol or nonoxynol phosphates, taurates, fatty taurides, sulfated monoglycerides, fatty acid amino polyoxyethylene sulfates and isothienates; or combinations thereof. Many additional anionic cleansing surfactants are described in McCUTCHEON'S DETERGENTS AND EMULSIFIERS, 1989 ANNUAL published by McCutcheon's Division MC Publishing Company, herein incorporated by reference. Usually, the anionic cleansing surfactant is present in the composition as a neutralized salt in the form of a sodium, potassium, lithium, ammonium, alkylammonium or hydroxyalkylammonium salt, wherein the alkyl moiety includes from one to about three carbon atoms.

[0025] Exemplary anionic cleansing surfactants that are useful in the composition and method of the present invention

include, but are not limited to, the ammonium, monoethanolamine, diethanolamine, triethanolamine, isopropylamine, sodium, potassium, lithium, or magnesium salts of lauryl sulfate, dodecylbenzenesulfonate, lauryl sulfosuccinate, lauryl ether sulfate, lauryl ether carboxylate, lauryl sarcosinate, cocomethyl tauride, and sulfosuccinate half ester amide; or combinations thereof. Also useful are the zwitterionic betaines, e.g., cocamidopropyl betaine, cocamidopropyl hydroxysultaine, and the like; and the anionic carboxylate cleansing detergents, such as C₁₁-C₁₅ Pareth-7 carboxylic acid, C₁₁-C₁₅ Pareth-9, C₁₁-C₁₅ Pareth-12, C₁₁-C₁₅ Pareth-20, C₁₁-C₁₅ Pareth-30, C₁₁-C₁₅ Pareth-40, C₁₁-C₂₁ Pareth-10, C₁₂-C₁₃ Pareth-5 carboxylic acid, C₁₂-C₁₅ Pareth-2 phosphate, C₁₂-C₁₅ Pareth-7 carboxylic acid, C₁₂-C₁₅ Pareth-9, C₁₂-C₁₅ Pareth-12, C₁₄-C₁₅ Pareth-13, C₂₂-C₂₄ Pareth-33, cocaminobutyric acid, cocaminopropionic acid, coceth-7 carboxylic acid, cocoamphodipropionic acid, coconut acid, deceth-7 carboxylic acid, hydrogenated coconut acid, hydrogenated menhaden acid, hydrogenated tallow acid, hydroxystearic acid, isostearic acid, lanolin acid, lauraminopropionic acid, laureth-5 carboxylic acid, laureth-10 carboxylic acid, lauroamphodipropionic acid, linoleic acid, linolenic acid, linseed acid, MEA-laureth-6-carboxylate, myristaminopropionic acid, palmitic acid, sodium C₁₂-C₁₅ Pareth-6 carboxylate, sodium C₁₂-C₁₅ Pareth-7 carboxylate, sodium ceteth-13 carboxylate, sodium isosteareth-6 carboxylate, sodium isosteareth-11 carboxylate, sodium laureth-13 carboxylate, sodium trideceth-7 carboxylate, sodium trideceth-12 carboxylate, trideceth-4 carboxylic acid, trideceth-7 carboxylate acid, trideceth-15 carboxylic acid, and trideceth-19 carboxylic acid.

[0026] The following low-irritation surfactants are particularly useful in formulating a "baby" shampoo having high performance in terms of foam level and cleansing while achieving exceptional mildness:

20 **ANIONICS:**

[0027]

Disodium Laureth Sulfosuccinate;
 Disodium Lauroamido MEA Sulfosuccinate;
 Disodium Ricinoleamido MEA Sulfosuccinate;
 Cetareth-25-Carboxylic Acid;
 Trideceth-7-Carboxylic Acid;
 Pareth-25-6-Carboxylic Acid;
 Trideceth-4-Carboxylic Acid;
 Trideceth-19-Carboxylic Acid;
 Sodium Trideceth-12-Carboxylate;
 Sodium Ceteth-13-Carboxylate;
 Laureth-5-Carboxylic Acid (SANDOPAN® LAB) ;
 Sodium Laureth-13-Carboxylate;
 Sodium Oleth-13-Carboxylate;
 Sodium Cetareth-5-Carboxylate;
 Sodium Cetareth-9-Carboxylate;
 Isosteareth-6-Carboxylic Acid; and
 Isosteareth-11-Carboxylic Acid.

NONIONICS:

[0028]

PEG 30 Glyceryl Mono Cocoate;
 PEG 78 Glyceryl Mono Cocoate;
 PEG 82 Glyceryl Mono Tallowate;
 PEG 200 Glyceryl Mono Tallowate; and
 PEG 20 Glyceryl Mono Tallowate.

AMPHOTERICS:


[0029]

Cocampho-Carboxyglycinate (VARION® 2C) ;
 Lauroampho-Carboxyglycinate (VARION® 2L) ;
 Cocamidopropyl Betaine; and

Cocamidopropyl Hydroxysultaine (VARION® CAS).

[0030] The polyethyleneimine(s) contained in the conditioning shampoos of the present invention generally have the formula $(\text{CH}_2\text{CH}_2\text{NH})_n$ wherein n has an average value of about 5 to about 2500. Specific examples of polyethyleneimines are PEI-7; PEI-15; PEI-30; PEI-45; PEI-1000; PEI-1500; and PEI-2500, wherein the integer following the PEI corresponds to the value of n in the formula above.

[0031] Surprisingly, protonated polyethyleneimines, together with the cationic Polyquaternium 32, provide excellent stability while achieving additional conditioning benefits from the protonated polyethyleneimine. The polyethyleneimines can be obtained protonated with any suitable acid at a pH below about 7.0, or the polyethyleneimines can be protonated in-situ (during mixing of the composition with an acid, e.g., citric acid) by adding sufficient acid that is free to protonate the polyethyleneimine. The molecular weight of the polyethyleneimine is not critical and can be any molecular weight commercially available, e.g., protonated polyethyleneimines available from BASF Corporation having a weight average molecular weight in the range of about 700 to about 70,000. The ability to provide a conditioning shampoo that has excellent conditioning benefits, as well as excellent foaming and stability, has been a long-felt need in the conditioning shampoo art. The conditioning shampoos of the present invention solve this long-felt need by including a protonated polyethyleneimine and a cationic acrylate/acrylamide copolymer, particularly Polyquaternium 32. The structure of protonated polyethyleneimine is as follows:



[0032] The preferred protonated polyethyleneimines have a ratio of primary:secondary:tertiary nitrogen atoms of about 1:2:1, respectively.

[0033] The emulsified conditioning shampoo of the present invention also includes a cationic conditioning agent, such as a SALCARE SC92 acrylate/acrylamide copolymer conditioning agent of Allied Colloids (POLYQUATERNIUM 32) as disclosed in our parent application Serial No. 07/719,818 filed June 24, 1991, hereby incorporated by reference.

[0034] To achieve the full advantage of the present invention, an optional foam booster, in an amount of about 0.1% to about 20% by weight of the composition, is included in the composition to aid in the formation of copious amount of foam. Suitable foam boosters include one or more of the following:

Capramide DEA	Disodium Isostearyl Sulfosuccinate
Cetearyl Alcohol	Hydrogenated Tallow Amine Oxide
Cetyl Alcohol	Hydroxyethyl Hydroxypropyl C ₁₂ -C ₁₅ Alkoxypropylamine Oxide
Cetyl Betaine	Hydroxyethyl Stearamide-MIPA
Cocamide	Isopropyl Stearate
Cocamide DEA	Isostearamidopropylamine Oxide
Cocamide MEA	Isostearamidopropyl Morpholine Oxide
Cocamide MIPA	Lauramide

EP 0 627 217 B1

(continued)

	Cocamidoethyl Betaine	Lauramide DEA
5	Cocamidopropylamine Oxide	Lauramide MEA
	Cocamidopropyl Betaine	Lauramide MIPA
	Cocamidopropyl Hydroxysultaine	Lauramidopropylamine Oxide
	Cocamine Oxide	Lauramidopropyl Betaine
10	Cocoamphodipropionic Acid	Lauramine Oxide
	Coco-Betaine	Lauryl Alcohol
	Coco-Morpholine Oxide	Lauryl Betaine
15	Coconut Alcohol	Lauryl Sultaine
	Coco/Oleamidopropyl Betaine.	Myristamide DEA
	Coco-Sultaine	Myristamide MEA
	Cocoyl Hydroxyethyl Imidazoline	Myristamide MIPA
20	Cocoyl Sarcosinamide DEA	Myristamidopropylamine Oxide
	DEA-Cocoamphodipropionate	Myristamidopropylamine Betaine
	DEA-Lauraminopropionate	Myristamine Oxide
25	Decylamine Oxide	Myristaminoproionic Acid
	Decyl Betaine	Myristyl Alcohol
	Dihydroxyethyl C ₈ -C ₁₀ Alkoxypropylamine Oxide	Myristyl Betaine
	Dihydroxyethyl C ₉ -C ₁₁ Alkoxypropylamine Oxide	Oleamidopropylamine Oxide
30	Dihydroxyethyl C ₁₂ -C ₁₅ Alkoxypropylamine Oxide	Oleamidopropyl Betaine
	Dihydroxyethyl Cocamine Oxide	Oleamidopropyl Hydroxysultaine
	Dihydroxyethyl Stearamine Oxide	Oleamine Oxide
35	Dihydroxyethyl Tallowamine Oxide	Oleyl Betaine
		Palmamide DEA

40	Palmamide MEA	PEG-5 Lauramide
	Palmamide MIPA	PEG-6 Lauramide
	Palmamidopropyl Betaine	PEG-3 Lauramine Oxide
45	Palmitamide DEA	Sodium Cocoamphoacetate
	Palmitamide MEA	Sodium Cocoamphopropionate
	Palmitamidopropylamine Oxide	Sodium Lauraminopropionate
	Palmitamidopropyl Betaine	Sodium Lauroamphopropionate
50	Palmitamine Oxide	Sodium Lauroyl Sarcosinate
	Palm Kernel Alcohol	Sodium Myristoamphoacetate
	Palm Kernelamide DEA	Sodium Myristoyl Sarcosinate
	Palm Kernelamide MEA	Stearyl Alcohol
55	Palm Kernelamide MIPA	TEA-Hydrogenated Tallow Glutamate
	Peanutamide MEA	TEA-Lauraminopropionate

(continued)

Peanutamide MIPA	TEA-Myristaminopropionate
PEG-6 Cocamide	Undecylenamide DEA
PEG-3 Lauramide	Undecylenamide MEA
	Undecylenamidopropylamine Oxide

[0035] One or more zwitterionic detergents, such as a betaine, in an amount of about 5% to about 25% by weight of the composition aids in stabilizing the composition but generally is not necessary to achieve a stable composition. Suitable betaines include, for example:

Betaine	Myristamidopropyl Betaine
Cetyl Betaine	Myristyl Betaine
Cocamidoethyl Betaine	Oleamidopropyl Betaine
Cocamidopropyl Betaine	Oleamidopropyl Hydroxysultaine
Cocamidopropyl Hydroxysultaine	Oleyl Betaine
Coco-Betaine	Palmamidopropyl Betaine
Coco/Oleamidopropyl Betaine	Palmitamidopropyl Betaine
Coco-Sultaine	Ricinoleamidopropyl Betaine
Decyl Betaine	Stearamidopropyl Betaine
Hydrogenated Tallow Betaine	Stearyl Betaine
Isostearamidopropyl Betaine	Tallowamidopropyl Betaine
Lauramidopropyl Betaine	Tallowamidopropyl Hydroxysultaine
Lauryl Betaine	Wheat Germamidopropyl Betaine
Lauryl Sultaine	

[0036] Other compounds useful for composition stabilization, in an amount of about 0.1% to about 10% by weight of the composition include any one or more of the following:

Acetylated Glycol Stearate	Maltodextrin
Aluminum Caprylate	Methoxy PEG-22/Dodecyl Glycol Copolymer
Aluminum Dilinoleate	Methylcellulose
Aluminum Distearate	Microcrystalline Cellulose
Aluminum Isostearates/Laurates/ Palmitates	Microcrystalline Wax
Aluminum Isostearates/Laurates/ Stearates	Montmorillonite
Aluminum Isostearates/Myristates	Myristyl Alcohol
Aluminum Isostearates/Palmitates	Ozokerite
Aluminum Isostearates/Stearates	Pectin
Aluminum Lanolate	PEG-2M
Aluminum Myristates/Palmitates	PEG-5M
Aluminum Stearate	PEG-7M
Aluminum Stearates	PEG-9M
Aluminum Tristearate	PEG-14M

EP 0 627 217 B1

(continued)

	Beeswax	PEG-20M
5	Bentonite	PEG-23M
	C ₉ -C ₁₁ Alcohols	PEG-45M
	C ₁₂ -C ₁₃ Alcohols	PEG-90M
	C ₁₂ -C ₁₅ Alcohols	PEG-115M
10	C ₁₂ -C ₁₆ Alcohols	PEG-22/Dodecyl Glycol Copolymer
	C ₁₄ -C ₁₅ Alcohols	PEG-45/Dodecyl Glycol Copolymer
	C ₁₅ -C ₁₈ Glycol	Polyacrylic Acid
15	Calcium Carrageenan	Polyethylene
	Calcium Stearate	Polyvinyl Acetate
	Carbomer 910	Potassium Alginate
	Carbomer 934	Potassium Carrageenan
20	Carbomer 934P	PVM/MA Copolymer
	Carbomer 940	PVP/VA Copolymer
	Carbomer 941	Saccharated Lime
25	Carboxymethyl Hydroxyethylcellulose	Sodium Acrylate/Vinyl Alcohol Copolymer
	Carboxymethyl Hydroxypropyl Guar	Sodium C ₄ -C ₁₂ Olefin/Maleic Acid Copolymer
	Carrageenan	Sodium Carboxymethyl Dextran
	Cellulose Gum	Sodium Carrageenan
30	Ceresin	Sodium Cellulose Sulfate
	Cetearyl Alcohol	Sodium Polymethacrylate
	Cetyl Alcohol	Sodium Polynaphthalene Sulfonate
35	Cholesterol	Sodium Polystyrene Sulfonate
	Coconut Alcohol	Stearyl Alcohol
	Ethylene/Acrylate Copolymer	Stearyl/Vinyl Ether/Maleic Anhydride Copolymer
	Ethylene/Vinyl Acetate Copolymer	Styrene/Maleic Anhydride Copolymer
40	Guar Gum	Synthetic Beeswax
	Hydroxybutyl Methylcellulose	Synthetic Wax
	Hydroxyethylcellulose	Tallow Alcohol
45	Hydroxyethyl Ethylcellulose	Tragacanth Gum
	Hydroxypropylcellulose	Tridecyl Alcohol
	Hydroxypropyl Guar	Xanthan Gum
	Hydroxypropyl Methylcellulose	
50	Isopropyl Ester of PVM/MA Copolymer	
	Karaya Gum	
	Lanolin	
55	Lanolin Alcohol	
	Lauryl Alcohol	
	Locust Bean Gum	

[0037] Other compounds useful for composition stabilization, in an amount of 0.1% to 55% by weight of the composition include a long chain (C₁₂-C₂₂) amine oxide.

[0038] Other common cosmetic components and additives that can be incorporated into the conditioning shampoos of the present invention, as long as the basic properties of conditioning, cleansing and high foam levels are not adversely affected include, for example, fragrances, dyes, hair colorants, opacifiers, pearlescing agents, dandruff control agents, hydrotropes, foam stabilizers, solubilizers, preservatives, water softening agents, acids, bases, buffers and the like. These optional components and additives usually will be present in weight percentages of less than about 2% each, and from about 5% to about 10% by weight in total.

[0039] The vehicle of the hair-treating composition is generally predominantly water, but organic solvents also can be used in order to help solubilize compounds that are not sufficiently soluble in water. Suitable solvents include the lower alcohols like ethyl alcohol and isopropyl alcohol; polyols like glycerol; glycols or glycol ethers, like 2-butoxyethanol, ethylene glycol, ethylene glycol monoethyl ether, propylene glycol and diethylene glycol monoethyl ether or monoethyl ether; and mixtures thereof. These non-aqueous solvents can be present in the hair-treating composition of the present invention in an amount from about 1% to 100% by weight and, in particular, from about 5% to about 50% by weight, relative to the total weight of the carrier vehicle in the composition.

[0040] The conditioning shampoos of the present invention also can be thickened, for example, with sodium alginate; guar gum; xanthan gum; gum arabic; cellulose derivatives such as methylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose and carboxymethylcellulose; and various polymeric thickeners, such as polyacrylic acid derivatives. These thickeners are present in an amount ranging from about 0.1% to about 5%, and preferably from about 0.25% to about 1%, by weight relative to the total weight of the composition.

[0041] If instability of the composition is a problem, the composition also can include a suspending agent for the conditioning agent or other water-insoluble material, in an amount of about 0.5% to about 10%, by total weight of the composition. Suitable suspending agents are for example, distearyl amate (distearyl phthalamic acid); fatty acid alkanolamides; esters of polyols and sugars; polyethylene glycols; the ethoxylated or propoxylated alkylphenols; ethoxylated or propoxylated fatty alcohols; and the condensation products of ethylene oxide with long chain amides. These suspending agents, as well as numerous others not cited herein, are well known in the art and are fully described in literature, such as McCUTCHEON'S DETERGENTS AND EMULSIFIERS, 1989 Annual, published by McCutcheon Division, MC Publishing Co.

[0042] The conditioning shampoos also can include other emulsifiers, inorganic salts, humectants and similar materials to provide esthetic properties and desirable physical properties to the composition. Generally, such optional ingredients are present in weight percentages ranging from about 0.1% to about 10% each, and from about 0.1% to about 20% in total, relative to the total weight of the composition.

[0043] For example, representative nonionic surfactants that can be included in the hair-treating composition of the present invention include esters of polyols and sugars; the polyethoxylated and/or polypropoxylated alkylphenols; and the condensation products of ethylene oxide with long chain amides. All these nonionic surfactants, as well as numerous others not cited here, are well known in the art and are fully described in the literature.

[0044] The compositions of the present invention can be relatively viscous dispersions that are stable to phase separation at a temperature of about 20°C to about 25°C for a period of time of at least 24 hours after preparation, and typically are stable to phase separation indefinitely at such temperatures. The compositions of the present invention usually are emulsions that are stable to phase separation at a temperature of about 25°C for a period of about 24 hours after preparation. The emulsions should demonstrate sufficient stability to phase separation at temperatures normally found in commercial product storage and shipping to remain unaffected for period of one year or more.

[0045] The following examples illustrate various conditioning shampoos made in accordance with the present invention:

EXAMPLES 1-3

[0046]

		1 WT.%	2 WT.%	3 WT.%
1.	Water	30.00	30.00	30.00
2.	Polyvinylpyrrolidone (stabilizing agent)	1.00	1.00	1.00
3.	Polyethyleneimine (Polymin G-35 BASF) wt. av. M.W - 1,700 as measured by GPC (conditioning and stability)	1.00	1.00	1.00

EP 0 627 217 B1

(continued)

		1 WT.%	2 WT.%	3 WT.%
5	4. Acrylate/acrylamide copolymer (Allied Colloids Salcare SC92) (50% in mineral oil)	1.00	1.00	1.00
	5. Citric Acid (pH adjustment)	1.80	1.80	1.80
10	6. Cocamide DEA (foam booster)	1.00	1.00	1.00
15	7. Surfactant blend (cleansing)	35.00	35.00	35.00
20	(a) ALES (1 mole ethoxylation) (8.0%)			
	(b) lauramide DEA (3.5%)			
	(c) ammonium xylene sulfonate (0.5%)			
	(d) water q.s.			
	8. Silicone blend 33 % SE 30 67% SF96-350 (conditioners)	0.65	0.65	0.65
25	9. Water	8.00	10.00	9.50
	10. Sodium lauryl sulfate (cleansing, foam)	15.00	15.00	15.00
30	11. Cocamidopropyl Hydroxysultaine (40%) (VARION® CAS) (amphoteric surfactant)	4.00	2.00	2.00
	12. Cocamide DEA (foam booster)	1.00	1.00	1.00
35	13. Kathon CG (preservative)	0.05	0.05	0.05
	14. Glydant (preservative)	0.20	0.20	0.20
	15. Fragrance	0.30	0.30	0.30
40	pH	4.2	4.4	4.2
	Viscosity	2 Pas	2,25 Pas	2,15 Pas

Mixing Procedure Examples 1-3

45 [0047] Add the polyvinylpyrrolidone (#2) to water (#1) with high agitation. Then add the polyethyleneimine (#3) and continue mixing. Then add the acrylate/acrylamide copolymer (#4) and mix until homogeneous (lump free). Then add the citric acid (#5), the cocamide DEA (#6), the surfactant blend (#7) and the silicone blend (#8). Into a separate container add the sodium lauryl sulfate (#10) to water (#9) and then add the contents of the separate container to the mixture of (#1) through (#8), with agitation. Then add the VARION® CAS (#13), the glydant (#14) and the fragrance (#15).

EXAMPLE 4

[0048]

55

		WT.%
1.	Water	59.05

EP 0 627 217 B1

(continued)

		WT. %
2.	Polyethyleneimine (Polymin G35, BASF) (conditioning and stability)	0.10
3.	Acrylate/acrylamide copolymer Allied Colloids D.P. 64297C (50% in mineral oil)	2.50
4.	SURFODONE® QSP (polylauryl pyrrolidone) polymeric emulsion stabilizer	0.50
5.	SANDOPAN® LA 8 (carboxylate surfactant)	7.00
6.	SURFINE® WNT A (carboxylate surfactant)	7.00
7.	KOH (50%)	0.60
8.	Cocamidopropyl Hydroxysultaine (40%) (VARION® CAS) (amphoteric surfactant)	15.00
9.	C ₁₆₋₁₈ amine oxide (40%) (SCHERCAMOX® CMA)	2.00
10.	GLUCAMATE® DOE 120* (thickener)	0.50
11.	Color, fragrance, preservative pH = 5.19 Viscosity (25°C) = 3 Pa s.	q.s.

* polyethylene glycol diester of methyl glucose and oleic acid with an average of 120 moles of ethylene oxide.

EXAMPLE 5

[0049]

		WT. %
1.	Water	19.50
2.	Polyethyleneimine (Polymin P, BASF) (conditioning and stability)	0.50
3.	Acrylate/acrylamide copolymer Allied Colloids D.P. 64297C (50% in mineral oil)	2.00
4.	Cocamidopropyl Hydroxysultaine (40%) (VARION® CAS) (amphoteric surfactant)	15.00
5.	Cocamide DEA (foam booster)	4.00
6.	SANDOPAN® LA 8 (carboxylate surfactant)	15.00
7.	Water	42.05
8.	KOH (50%)	1.40
9.	Perfume	0.30
10.	Glydant	0.20
11.	Kathon CG pH - 5.0 Viscosity (25°C) = 2 Pa s	0.05

Mixing Procedure Example 5:

5 [0050] Add the sultaine (#4), the polyethyleneimine (#2), and the acrylate/acrylamide copolymer (#3) to water (#1) with mixing (1 hour). Then add the Cocamide DEA (#5) and mix until homogeneous (lump free). Into a separate container, add the SANDOPAN® carboxylate surfactant (#6) to water (#7) followed by the KOH (#8) addition and then mix all items (#1) through (#8). The perfume (#9), glydant (#10) and Kathon CG (#11) then are added.

EXAMPLE 6

10 [0051]

		WT. %
15	1. Water	35.50
	2. Polyethyleneimine (POLYMIN WATERFREE, BASF) (conditioning and stability)	0.50
20	3. SURFADONE® QSP (polylauryl pyrrolidone) polymeric emulsion stabilizer	0.60
	4. Acrylate/acrylamide copolymer Allied Colloids D.P. 65780E (50% mineral oil)	3.00
25	5. Surfactant Blend: (a) ALES (1 mole ethoxylation) (4.5%) (b) lauramide DEA (2.5%) (c) ammonium xylene sulfonate (0.6%) (d) water q.s.	20.00
30	6. KOH (50%)	0.50
35	7. SANDOPAN® LA 8 (carboxylate surfactant)	15.00
	8. KOH	1.00
	9. Cocamidopropyl Hydroxysultaine (40%) (VARION® CAS) (amphoteric surfactant)	10.00
40	10. Perfume	0.50
	11. Glydant	0.20
	12. Kathon CG	0.05
	13. Cold pearl mix	3.00
45	14. Cocamide DEA FO (foam booster)	3.00
	15. Water, color	7.15

Mixing Procedure Example 6:

50 [0052]

Add (#2) and (#3) in (#1), mix until no lumps, then add (#4) and mix until a soft gel is formed in about 40 minutes, avoid aeration;

Add (#5), mix for 5 minutes (precipitation is observed);

55 Add (#6) and (#7) and mix for 5 minutes;

Add (#8), mix until product is uniform (pH 4.5);

Add (#9), gradually, add (#10), (#11), (#12), (#13), (#14) and (#15).

pH 4.5

ADD KOH (50%) - 0.6% to pH 5.13**FINAL** pH = 5.13

5 Viscosity (25°C) = 6,3 Pa s

[0053] To show that the compositions of the present invention are compatible with a relatively low percentage of sodium lauryl ether (1 mole of ethoxylation) sulfate (SLES), Example 7 incorporates 5% by weight of sodium lauryl ether sulfate (1 mole of ethoxylation) (SLES) together with about 15% of other, mild carboxylate anionic surfactants.

EXAMPLE 7**[0054]**

15

20

25

30

35

40

		WT. %
1.	Water, Soft	20.72
2.	Polyethyleneimine (POLYMIN PS, BASF) (conditioning and stability)	2.00
3.	SURFODONE® QSP (polylauryl pyrrolidone) polymeric emulsion stabilizer	0.38
4.	Acrylate/acrylamide copolymer Allied Colloids D.P. 65789E (50% in mineral oil)	1.90
5.	Water	22.75
6.	SANDOPAN® LA 8 (90%) (carboxylate surfactant)	15.00
7.	KOH (50%)	1.50
8.	SLES (25%)	20.00
9.	Cocamidopropyl Hydroxysultaine (40%) (VARION® 2L) (amphoteric surfactant)	10.00
10.	Cocamide DEA (foam stabilizer)	3.00
11.	Perfume	0.50
12.	Kathon CG	0.05
13.	Clydant	0.20
14.	C ₁₆₋₁₈ amide oxide (40%) SHERCOMOX® CMA pH = 6.0 Viscosity (25°C) = 8,5 Pa s	2.00

Mixing Procedure Example 7:

45

[0055] Add (#2) to (#3) to (#1), mix well and add (#4), mix until soft gel is formed.**[0056]** Add remaining ingredients. Allow 5 minutes of mixing for each addition.

50

Claims

55

1. A mild conditioning shampoo capable of thoroughly cleansing and conditioning hair while maintaining a high foam level comprising water, an anionic cleansing surfactant in an amount of about 5% to about 65% by weight; a cationic polyethyleneimine in an amount of about 0.01% to about 4% by weight; and a cationic oil-soluble, water-dispersible, cross-linked acrylate/acrylamide copolymer conditioning agent in an amount of about 0.1% to about 20% by weight.
2. The composition of Claim 1, wherein the anionic cleansing surfactant is a carboxylate surfactant.

3. The composition of Claims 1 or 2 further including a long chain (C₁₂-C₂₂) amine oxide emulsion stabilizer in an amount of about 0.1% to about 55 based on the weight of the composition.
- 5 4. The composition of anyone of Claims 1 to 3 having a pH of about 4 to about 7.
5. The composition of anyone of Claims 1 to 4, wherein the surfactant includes a viscosity increasing agent in an amount of about 0.1% to about 10% by weight capable of raising the viscosity of the composition to at least about 3 Pa s.
- 10 6. The composition of anyone of Claims 1 to 5 further including a zwitterionic detergent in an amount of about 5 % to about 25 % by weight of the composition, preferably a betaine surfactant, such as cocamidopropyl hydroxysultaine.
- 15 7. The composition of anyone of Claims 1 to 6, further including a silicone conditioning agent in an amount of about 0.1% to about 10% by weight of the composition.
8. The composition of anyone of Claims 1 to 7, wherein the composition includes less than about 9% by weight of a strong anionic surfactant which is preferably selected from the group consisting of a long chain (C₁₂-C₂₂) alkyl sulfate; long chain (C₁₂-C₂₂) alkyl ether sulfate; a long chain (C₁₂-C₂₂) alkyl sulfonate; and a long chain (C₁₂-C₂₂) alkyl ether sulfonate.
- 20 9. The composition of anyone of Claims 1 to 8, wherein the polyethyleneimine has a cationic polymer charge density of at least about 10 milliequivalents per gram, preferably in the range of about 15 to about 20 milliequivalents per gram.
- 25 10. A method of cleansing and conditioning hair, simultaneously, while maintaining a substantial quantity of foam and excellent cleansing in a mild shampoo composition comprising contacting the hair with a mild conditioning shampoo as defined in anyone of Claims 1 to 9.

30

Patentansprüche

- 35 1. Mildes konditionierendes Shampoo, das Haar unter Halten eines hohen Schaumniveaus gründlich reinigen und konditionieren kann, umfassend Wasser, ein anionisches reinigendes Tensid in einer Menge von etwa 5 Prozent bis etwa 65 Gewichtsprozent; ein kationisches Polyethylenimin in einer Menge von etwa 0,01 Prozent bis etwa 4 Gewichtsprozent; und ein kationisches, in Öllösliches, in Wasser dispergierbares, vernetztes Acrylat/Acrylamid-Copolymer-Konditionierungsmittel in einer Menge von etwa 0,1 Prozent bis etwa 20 Gewichtsprozent.
- 40 2. Zusammensetzung nach Anspruch 1, wobei das anionische reinigende Tensid ein Carboxylattensid darstellt.
3. Zusammensetzung nach Ansprüchen 1 oder 2, die weiterhin einen langkettigen (C₁₂-C₂₂)-Aminoxid-Emulsionsstabilisator in einer Menge von etwa 0,1 Prozent bis etwa 55 Gewichtsprozent, bezogen auf das Gewicht der Zusammensetzung, einschließt.
- 45 4. Zusammensetzung nach einem der Ansprüche 1 bis 3 mit einem pH-Wert von etwa 4 bis etwa 7.
5. Zusammensetzung nach einem der Ansprüche 1 bis 4, wobei das Tensid ein Viskositäts-erhöhendes Mittel in einer Menge von etwa 0,1 Prozent bis etwa 10 Gewichtsprozent einschließt, das die Viskosität der Zusammensetzung auf mindestens etwa 3 Pa.s steigern kann.
- 50 6. Zusammensetzung nach einem der Ansprüche 1 bis 5, die weiterhin ein zwitterionisches Waschmittel in einer Menge von etwa 5 Prozent bis etwa 25 Gewichtsprozent der Zusammensetzung, vorzugsweise ein Betaintensid, wie Cocamidopropylhydroxysultain, einschließt.
- 55 7. Zusammensetzung nach einem der Ansprüche 1 bis 6, die weiterhin ein konditionierendes Silikonmittel in einer Menge von etwa 0,1 Prozent bis etwa 10 Gewichtsprozent der Zusammensetzung einschließt.
8. Zusammensetzung nach einem der Ansprüche 1 bis 7, wobei die Zusammensetzung weniger als etwa 9 Gewichts-

prozent eines starken anionischen Tensids einschließt, das vorzugsweise aus der Gruppe, bestehend aus einem langkettigen (C₁₂-C₂₂)-Alkylsulfat; langkettigen (C₁₂-C₂₂)-Alkylethersulfat; einem langkettigen (C₁₂-C₂₂)-Alkylsulfonat; und einem langkettigen (C₁₂-C₂₂)-Alkylethersulfonat, ausgewählt ist.

- 5 9. Zusammensetzung nach einem der Ansprüche 1 bis 8, wobei das Polyethylenimin eine kationische Polymerladungsdichte von mindestens etwa 10 Milliäquivalenten pro Gramm, vorzugsweise im Bereich von etwa 15 bis etwa 20 Milliäquivalenten pro Gramm, aufweist.
- 10 10. Verfahren zum gleichzeitigen Reinigen und Konditionieren von Haar, während eine wesentliche Menge Schaum und ausgezeichnetes Reinigen in einer milden Shampoозusammensetzung beibehalten wird, umfassend in Kontakt bringen des Haars mit einem, wie in einem der Ansprüche 1 bis 9 definierten, milden konditionierenden Shampoo.

15 **Revendications**

- 20 1. Shampoing de conditionnement doux capable de nettoyer et de conditionner les cheveux à fond tout en maintenant un niveau de mousse élevé, comprenant de l'eau, un tensioactif de nettoyage anionique dans une quantité allant d'environ 5 % à environ 65 % en poids ; un polyéthylèneimine cationique dans une quantité d'environ 0,01 % à environ 4 % en poids ; et un agent de conditionnement copolymère cationique acrylate/maléate réticulé, dispersible dans l'eau et soluble dans l'huile dans une quantité allant d'environ 0,1 % à environ 20 % en poids.
- 25 2. Composition selon la revendication 1, dans laquelle le tensioactif de nettoyage anionique est un tensioactif carboxylate.
- 30 3. Composition selon la revendication 1 ou 2, incluant en outre un stabilisateur d'émulsion d'oxyde d'amine à longue chaîne (C₁₂ - C₂₂) dans une quantité allant d'environ 0,1 % à environ 55, sur la base du poids de la composition.
4. Composition selon l'une des revendications 1 à 3, ayant un pH d'environ 4 à environ 7.
- 35 5. Composition selon l'une quelconque des revendications 1 à 4, dans laquelle le tensioactif inclut un agent d'augmentation de la viscosité dans une quantité allant d'environ 0,1 % à environ 10 % en poids, capable d'augmenter la viscosité de la composition à un niveau d'au moins environ 3 Pas.
- 40 6. Composition selon l'une quelconque des revendications 1 à 5, incluant en outre un détergent zwitterionique dans une quantité allant d'environ 5 % à environ 25 % en poids de la composition, de préférence un tensioactif bétaine tel que de la cocamidopropyl hydroxysultaine.
- 45 7. Composition selon l'une quelconque des revendications 1 à 6, incluant en outre un agent de conditionnement silicone dans une quantité allant d'environ 0,1 % à environ 10 % en poids de la composition.
8. Composition selon l'une quelconque des revendications 1 à 7, dans laquelle la composition inclut moins d'environ 9 % en poids d'un tensioactif fortement anionique qui est de préférence sélectionné à partir du groupe constitué d'un alkyl sulfate à longue chaîne (C₁₂-C₂₂), d'un alkyéther sulfate à longue chaîne (C₁₂ - C₂₂), d'un alkyl éther sulfonate à longue chaîne (C₁₂-C₂₂).
- 50 9. Composition selon l'une quelconque des revendications 1 à 8, dans laquelle le polyéthylèneimine a une densité de charge cationique d'au moins environ 10 milliéquivalents par gramme, de préférence dans la gamme allant d'environ 15 à environ 20 milliéquivalents par gramme.
- 55 10. Procédé de nettoyage et de conditionnement des cheveux, de manière simultanée, tout en maintenant une quantité substantielle de mousse est un excellent nettoyage dans une composition de shampoing doux et comprenant le fait de mettre en contact les cheveux avec un shampoing de conditionnement doux tel que défini dans l'une quelconque des revendications 1 à 9.

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)